

Relative humidity dependence of light absorption by mineral dust after long-range atmospheric transport from the Sahara

Daniel A. Lack,^{1,2} Patricia K. Quinn,³ Paola Massoli,^{1,2,4} Timothy S. Bates,³ Derek Coffman,³ David S. Covert,⁵ Berko Sierau,^{5,6} Sara Tucker,¹ Tahllee Baynard,^{1,2} Edward Lovejoy,¹ Daniel M. Murphy,¹ and A. R. Ravishankara¹

Received 15 September 2009; revised 6 November 2009; accepted 17 November 2009; published 19 December 2009.

[1] The relative humidity (RH) dependence of light absorption for a Saharan dust-dominated air mass transported to the Gulf of Mexico was measured during the 2006 TexAQS/GoMACCS study using a photo-acoustic absorption spectrometer (PAS). Aerosol absorption was measured at low (25%) and high (73%) RH indicating a $1.5(\pm 0.3)$ absorption enhancement [$f(\text{RH}_{\text{Abs}})$] under high RH conditions. $f(\text{RH}_{\text{Abs}})$ estimates, based on air-mass physical and optical properties and Mie theory modeling, were between 1.2–1.4. Reasons for differences between the measured and modeled $f(\text{RH}_{\text{Abs}})$ are discussed. The mass absorption coefficient of the long-range transported dust was calculated to be $0.04(\pm 0.02) \text{ m}^2\text{g}^{-1}$. **Citation:** Lack, D. A., et al. (2009), Relative humidity dependence of light absorption by mineral dust after long-range atmospheric transport from the Sahara, *Geophys. Res. Lett.*, 36, L24805, doi:10.1029/2009GL041002.

1. Introduction

[2] The optical properties of atmospheric particles can be dependent on the RH of the atmosphere in which they exist. If particles absorb sufficient water, surface area and scattered light increases. Parameterizations of this RH effect are included in climate models to better predict the radiative forcing of hygroscopic scattering aerosol [e.g., *Schmidt et al.*, 2006]. The ratios between wet and dry scattering or extinction [$f(\text{RH}_{\text{Sca}})$, $f(\text{RH}_{\text{Ext}})$] have been measured using nephelometry [*Carrico et al.*, 2003; *Li-Jones et al.*, 1998] and cavity ring down aerosol extinction spectroscopy (CRD-AES) [*Baynard et al.*, 2007] respectively. While $f(\text{RH}_{\text{Sca}})$ is relatively well understood from theoretical, laboratory and field investigations, quantification of $f(\text{RH})$ on particle light absorption [$f(\text{RH}_{\text{Abs}})$] is based on modeling or limited laboratory measurements [e.g., *Lewis et al.*, 2009; *Zhang et al.*, 2008]. Light absorption can be enhanced as a particle is coated in a scattering layer that focuses the

radiation onto the core [*Ackerman and Toon*, 1981; *Fuller*, 1995]. *Bond et al.* [2006] calculated absorption enhancements ($E_{\text{Abs}} = \text{Abs}_{\text{Coated}}/\text{Abs}_{\text{Non-Coated}}$) for typical atmospheric distributions of black carbon (BC) and semi-volatile coatings and found an E_{Abs} of 1.5 likely. The lack of measurements of $f(\text{RH}_{\text{Abs}})$ is due to instrument limitations; for example filter-based and photo-acoustic techniques show filter sensitivity to RH and signal loss due to evaporation of semi-volatile material respectively [*Lewis et al.*, 2009; *Murphy*, 2009; *Schmid et al.*, 2006]. Ambient measurement of $f(\text{RH}_{\text{Abs}})$ using the difference of extinction and scattering is prohibited by the uncertainties of the derived absorption.

[3] Many studies have calculated the radiative impacts of dust and found that the atmospheric burden of 1000–5000 Tgyr⁻¹ [*Heintzenberg*, 2009] contributes a substantial climate influence from globally averaged cooling to altered precipitation patterns [*Rosenfeld and Rudich*, 2001; *Shell and Somerville*, 2007; *Solmon et al.*, 2008]. Saharan dust is an extensively studied dust source, can absorb 1 to 10% of the radiation it encounters (i.e. single scattering albedo (SSA) of ~ 0.90 at mid visible wavelengths) and has variable wavelength absorption [*Müller et al.*, 2009; *Solmon et al.*, 2008]. The size and optical properties of lofted dust can change as transport occurs; e.g. the SSA can increase and chemical composition change as it becomes internally mixed with hygroscopic materials such as sulfate [*Andreae et al.*, 1986; *Levin et al.*, 1996], sea salt [*Levin et al.*, 2005] and ammonium nitrate [*Noble and Prather*, 1996]. Oxidation of SO₂ to sulfate on dust particles also occurs [*Usher et al.*, 2003]. Up to 80% of transported dust can be internally mixed with other material [*Andreae et al.*, 1986; *Trochkin et al.*, 2003; *Zhang et al.*, 2003] and suspension of hygroscopic dust into the atmosphere can occur [*Rudich et al.*, 2002]. The $f(\text{RH}_{\text{Sca}})$ (and size growth due to RH changes, $g(\text{RH}_{\text{Sca}})$) of mineral dust aerosol have been measured to be between 1.1 and 1.7 [*Carrico et al.*, 2003; *Howell et al.*, 2006; *Kaaden et al.*, 2009; *Li-Jones et al.*, 1998], with lower values usually corresponding to dust measured close to the source region. *Carrico et al.* [2003] concluded that, due to the high ratio of external to internal mixing, there was no appreciable change to the radiative properties due to dust becoming internally mixed with hygroscopic material, rather the RH response in scattering was dominated by the externally mixed aerosol. *Bates et al.* [2006] reviewed the radiative impact of RH on dust aerosol and concluded that the effect was negligible and not necessary to accurately model the direct radiative effect. To date, there have not been any field measurements of $f(\text{RH}_{\text{Abs}})$ of dust or BC.

¹Earth Systems Research Laboratory, NOAA, Boulder, Colorado, USA.

²Cooperative Institute for Research of the Environmental Sciences, University of Colorado, Boulder, Colorado, USA.

³Pacific Marine Environment Laboratory, NOAA, Seattle, Washington, USA.

⁴Now at Aerodyne Research, Inc., Billerica, Massachusetts, USA.

⁵Atmospheric Sciences Department, University of Washington, Seattle, Washington, USA.

⁶Now at Institute for Atmospheric and Climate Science, Swiss Federal Institute of Technology, Zurich, Switzerland.




Figure 1. (a) Absorption wavelength dependence (\dot{A}_{Abs}) versus dust fraction of total mass for all data. Major dust events shown in color, other data in black. (b) Chemical composition of particles sampled during the dust event of interest. (c) Average volumetric size distribution of the particles during the event of interest. (d) Flexpart footprint emission sensitivity corresponding to the 29 August 2006 sampling period.

Here we present $f(\text{RH}_{\text{Abs}})$ measured for dust transported from the Saharan region of Africa to the Gulf of Mexico.

2. Field Observations

[4] Chemical, optical and physical properties of $\leq 10 \mu\text{m}$ aerodynamic diameter ($d_p < 10 \mu\text{m}$) aerosol were measured during the 2006 TexAQS/GoMACCS study onboard the NOAA R/V *Ronald H. Brown* (RHB) and are described by *Quinn and Bates* [2005] and *Bates et al.* [2008]. Mass concentrations ($\pm 11\%$) of particulate sulfate, nitrate, ammonium, organic matter (POM), EC and inorganic oxidized material (IOM) (dust proxy) were measured using impactor filters over 8–12 hour periods [*Quinn and Bates*, 2005]. Aerosol extinction ($\pm 1\%$ @ $\sim 25\%$ RH) and $f(\text{RH}_{\text{Ext}})$ ($\pm 5\%$ @ 75% RH) were measured at 532 nm using CRD-AES [*Baynard et al.*, 2007]. Aerosol absorption ($\pm 5\%$ @ $\sim 25\%$ RH) was measured at 532 nm using PAS [*Lack et al.*, 2006] and at 467 nm, 530 nm and 660 nm using a particle soot absorption photometer (PSAP, $\pm \sim 30\%$) [*Lack et al.*, 2008]. (Certain commercial equipment is identified in this article to adequately specify the experimental procedure. Such identification does not imply recognition or endorsement by NOAA nor does it imply that the equipment is necessarily the best available for the purpose.) During a dust-dominated period, the PAS was re-configured to sample at high RH ($\sim 73\%$). RH was measured using a Vaisala HMP50 probe with an uncertainty of $\pm 3\%$. The sample was passed through a wet permeable membrane (max. RH 90%) and the RH reduced by adjusting sample flow rate. Waterproof microphones (Knowles Acoustics WP3502) were used during the RH trial and the resonant frequency recalibrated at RH changes to remove instrument RH

dependence. Data collection for the CRD-AES and PAS was interrupted before and after the RH trial (and during for the CRD-AES) due to the instrument plumbing changes required to elevate the RH.

3. Results

[5] The $f(\text{RH}_{\text{Abs}})$ of dust-dominated particles was measured during a major dust ‘event’ (27–30th August 2006). Dust was identified as the dominant PM constituent using a combination of measurements: Figure 1a shows the wavelength dependence for absorption (\dot{A}_{Abs}), for the entire field campaign ranging from 0.5–6 with larger \dot{A}_{Abs} usually coinciding with the enhanced dust mass. Dust shows enhanced absorption at shorter wavelengths, resulting in a \dot{A}_{Abs} much greater than 1 [*Linke et al.*, 2006]. Across the event of interest dust was found to contribute 70% of $d_p < 10 \mu\text{m}$ particle mass ($41.4 \mu\text{g m}^{-3}$) (Figure 1b) and the aerosol had a volume median aerodynamic diameter of $2.1 \mu\text{m}$ (Figure 1c). Particle backscatter and vertical turbulence profiles measured using Doppler LIDAR [*Tucker et al.*, 2009] showed layers at 1–1.5 km mixing down to the surface around 1200 UTC 27th of August. Footprint emission sensitivities from the Flexpart Lagrangian dispersion model [*Stohl et al.*, 2005] indicate a North African source for the air-mass 10–15 days prior to measurement in Galveston Bay, Texas (Figure 1d). Average SSA of the event was $0.95(\pm 0.01)$ and average extinction and absorption coefficients were 50 and 2.5 Mm^{-1} respectively (@ 532 nm). Average $f(\text{RH}_{\text{Ext}})$ dropped from $\sim 2.4(\pm 0.3)$ prior to the event to $\sim 1.9(\pm 0.3)$ during the event (at 73% RH), higher than the expected hygroscopicity for pure dust. Hence, the following results are due to a particle ensemble




Figure 2. (a) PAS absorption (grey symbols), PSAP absorption (black line) and RH (grey line) time series. (b) Representation of the RH dependence of absorption of Saharan dust-dominated particles.

of mixed composition rather than pure dust. Over a 1-hour period on the 29th August 2006 the PAS was configured to measure dry ($\sim 25\%$) and high RH ($\sim 73 \pm 3\%$ RH) absorption. A difference in the absorption between RH levels was measured (Figure 2a). Dry absorption was monitored using a PSAP (Figure 2a, black line) and shown to be stable during the high RH trial (although PSAP $>$ PAS by $\sim 1 \text{ Mm}^{-1}$). Absorption data (1 sec.) 10–15min. either side of the RH change were used to produce absorption histograms. Gaussian curves were fit to each histogram and an $f(\text{RH}_{\text{Abs}})$ of $\sim 1.50(\pm 0.34)$ was calculated from the ratio of the mean values of the Gaussian distributions (Figure 2b). A 2-sample t-test was used to calculate the standard deviation in the $f(\text{RH}_{\text{Abs}})$.

4. Discussion

4.1. Photo-Acoustics

[6] The photo-acoustic technique provides a lower limit on absorption due to potential PAS signal reductions introduced by evaporation of water on the particle (at high RH) and internal heat losses at large particle sizes. Utilizing the work of *Murphy* [2009] we calculate, for the size and assumed optical properties of the dust in this study, that a 7–30% reduction in $f(\text{RH}_{\text{Abs}})$ can be expected due to the above mentioned effects (see auxiliary material) depending on the assumption of water accommodation coefficient.¹

4.2. Simulations of $f(\text{RH}_{\text{Abs}})$

[7] Using core-shell Mie theory [*Bohren and Huffman*, 1983] and assumed particle properties we show that an

$f(\text{RH}_{\text{Abs}})$ of up to 1.4 could be expected for the dust event studied. Dust particles are rarely spherical and assuming sphericity (to simplify the calculations) tends to over estimate scattering and absorption (depending on non-spherical shape) [*Bohren and Huffman*, 1983]. As the particles are coated they will become more spherical and so our calculations will underestimate $f(\text{RH}_{\text{Abs}})$. We use optical properties representative of lofted African dust for Summer/Autumn [*Israelevich et al.*, 2003] (refractive index [RI] of $1.45 - 0.015i$) and assume an aqueous sulfate or NaCl coating (RI of $1.4 + 0.0i$) [*Tang and Munkjelwitz*, 1994]. We calculate that an average 25% diameter growth of $2.1 \mu\text{m}$ dust particles is required to produce the measured $f(\text{RH}_{\text{Ext}})$ of 1.9. This 25% growth is used as a center point in growth for further modeling. Next we calculate the theoretical $f(\text{RH}_{\text{Abs}})$ using Mie theory, the average measured size distribution (Figure 1c), a real RI for dust of 1.45, coat RI of $1.4 + 0.0i$ and a range of imaginary RI for dust from literature [*Balkanski et al.*, 2007; *Israelevich et al.*, 2003; *Müller et al.*, 2009]. The size distribution is grown by the fraction on the y-axis of Figure 3 which shows that $f(\text{RH}_{\text{Abs}})$ of 1.2–1.4 would be common across a range of particle growth and imaginary RI, even at the lower imaginary RI for dust measured in the SAMUM experiment [*Müller et al.*, 2009]. The measured $f(\text{RH}_{\text{Abs}})$ of 1.5 (black line, Figure 3a) is unlikely for the assumed properties if one considers that the 25% growth in diameter calculated above (grey line, Figure 3a) would be for a full internal mixture. If




Figure 3. (a) Modeled $f(\text{RH}_{\text{Abs}})$ for the dust-dominated particles encountered in this study (Figure 1c). See text for model assumptions. Dashed line: $f(\text{RH}_{\text{Abs}})$ of an internal mixture for this study. Black symbol: $f(\text{RH}_{\text{Abs}})$ for a $\sim 50\%$ internal mixture with dust RI from literature. Black line: $f(\text{RH}_{\text{Abs}})$ of 1.5. (b) SSA reduction expected by including $f(\text{RH}_{\text{Abs}})$ effect relative to humidification of dust-dominated particles without $f(\text{RH}_{\text{Abs}})$.

¹Auxiliary materials are available in the HTML. doi:10.1029/2009GL041002.

we assume a diameter growth of 15% (a growth approximating $\sim 50\%$ internal mixture [Trochkin *et al.*, 2003]) an $f(\text{RH}_{\text{Abs}})$ of 1.25–1.30 is modeled (black dot, Figure 3a). Given that studies have reported 5 to 80% internal mixing [Andreae *et al.*, 1986; Zhang *et al.*, 2003], the $f(\text{RH}_{\text{Abs}})$ expected for dust-dominated particles may range from 1 to 1.4. Modeling also showed that the $f(\text{RH}_{\text{Abs}})$ will be larger if the absorbing component of the dust is concentrated (i.e., higher imaginary RI) and mixed with non-absorbing quartz (not shown here).

4.3. Mass Absorption Coefficient

[8] The mass absorption coefficient (MAC) for dust is calculated to be $0.04(\pm 0.02) \text{ m}^2 \text{ g}^{-1}$ (532 nm). We estimate that the $0.1(\pm 0.1) \mu\text{g m}^{-3}$ EC (Figure 1b) contributed 0.75 Mm^{-1} of the total $2.5(\pm 0.2) \text{ Mm}^{-1}$ absorption (assuming EC MAC of $7.5 \text{ m}^2 \text{ g}^{-1}$). Therefore the absorption of $37.4(\pm 5.0) \mu\text{g m}^{-3}$ dust was $1.75(\pm 0.75) \text{ Mm}^{-1}$. The calculated MAC for the long-range transported dust was therefore $0.04(\pm 0.02) \text{ m}^2 \text{ g}^{-1}$; similar to recent studies [Yang *et al.*, 2009]. The MAC used for EC is usually applied to fresh EC emissions and can reasonably increase by 50% when coated by scattering material [Bond *et al.*, 2006]. A 50% increase in absorption by EC would contribute $\sim 0.40 \text{ Mm}^{-1}$ to the $f(\text{RH}_{\text{Abs}})$ shown in Figure 2b, reducing the discrepancy between measured and modeled $f(\text{RH}_{\text{Abs}})$ identified above.

4.4. Radiative Impacts

[9] We have shown that water uptake by dust affects both scattering and absorption and thus SSA. SSA is an important factor that influences the global radiation balance [Balkanski *et al.*, 2007] and predicted precipitation patterns over Western Africa [Solmon *et al.*, 2008]. Carrico *et al.* [2003], Howell *et al.* [2006] and McNaughton *et al.* [2009] concluded that RH-induced scattering changes within mixed dust/pollution aerosol would be negligible due to the dominance of external mixing (i.e., interaction of dust and pollution would be minimal) and a cancellation of scattering due to redistribution of hygroscopic material from the fine to coarse modes when internal mixing occurred. Those studies did not consider an $f(\text{RH}_{\text{Abs}})$ effect. To gain an insight to this the impact of RH on the SSA of the dust-dominated particles studied, we incorporated the $f(\text{RH}_{\text{Abs}})$ effect into Mie theory. Figure 3b shows that the SSA expected for humidified dust-dominated aerosol with the $f(\text{RH}_{\text{Abs}})$ effect can be reduced from 0.01–0.03 compared to the SSA when $f(\text{RH}_{\text{Abs}})$ is not considered. This reduction is for reasonable growth conditions and does not consider optical changes due to redistribution of externally mixed material, which will complicate the impact of the $f(\text{RH}_{\text{Abs}})$ effect. This result is for long-range transported, well-aged dust and likely represents the end of a continuum beginning at the dust source (i.e., these results may not be as significant for less aged dust).

5. Summary

[10] Photo-acoustic absorption spectroscopy was used in the field to measure an absorption enhancement from water uptake [$f(\text{RH}_{\text{Abs}})$] of $1.5(\pm 0.3)$ (RH $\sim 73\%$) for Saharan dust-dominated aerosol. Simulations of the $f(\text{RH}_{\text{Abs}})$ for the aerosol encountered showed $f(\text{RH}_{\text{Abs}})$ of 1.2–1.4 is possible

given assumed particle properties and that SSA can be reduced by 0.01–0.03. A small amount of strongly absorbing EC, or more absorbing dust mixed with non-absorbing quartz (rather than uniformly mixed absorbing dust) may resolve the difference in modeled and observed $f(\text{RH}_{\text{Abs}})$. The observed $f(\text{RH}_{\text{Abs}})$ shows that PAS is capable of measuring an $f(\text{RH}_{\text{Abs}})$ signal and that dust aerosol can exhibit $f(\text{RH}_{\text{Abs}})$ due to water uptake by internally mixed hygroscopic material. It is recognized that this analysis is based on a small subset of data however the unique insights gained show cause for further investigation.

[11] **Acknowledgments.** Funded in part by NOAA's Climate Program. We thank Chris Cappa for useful comments and the crew of the Ronald H. Brown.

References

- Ackerman, T. P., and O. B. Toon (1981), Absorption of visible radiation in atmosphere containing mixtures of absorbing and non-absorbing particles, *Appl. Opt.*, *20*(20), 3661, doi:10.1364/AO.20.003661.
- Andreae, M. O., et al. (1986), Internal mixture of sea salt, silicates, and excess sulfate in marine aerosols, *Science*, *232*(4758), 1620–1623, doi:10.1126/science.232.4758.1620.
- Balkanski, Y., et al. (2007), Reevaluation of mineral aerosol radiative forcings suggests a better agreement with satellite and AERONET data, *Atmos. Chem. Phys.*, *7*(1), 81–95.
- Bates, T. S., et al. (2006), Aerosol direct radiative effects over the northwest Atlantic, northwest Pacific, and north Indian oceans: Estimates based on in-situ chemical and optical measurements and chemical transport modeling, *Atmos. Chem. Phys.*, *6*(6), 1657–1732.
- Bates, T. S., et al. (2008), Boundary layer aerosol chemistry during TexAQS/GoMACCS 2006: Insights into aerosol sources and transformation processes, *J. Geophys. Res.*, *113*, D00F01, doi:10.1029/2008JD010023.
- Baynard, T., et al. (2007), Direct measurement of atmospheric aerosol optical extinction using cavity ring-down spectroscopy: Field instrument development, *Aerosol Sci. Technol.*, *41*(4), 447–462, doi:10.1080/02786820701222801.
- Bohren, C. F., and D. R. Huffman (1983), *Absorption and Scattering of Light by Small Particles*, 530 pp., John Wiley, New York.
- Bond, T. C., G. Habib, and R. W. Bergstrom (2006), Limitations in the enhancement of visible light absorption due to mixing state, *J. Geophys. Res.*, *111*, D20211, doi:10.1029/2006JD007315.
- Carrico, C. M., et al. (2003), Mixtures of pollution, dust, sea salt, and volcanic aerosol during ACE-Asia: Radiative properties as a function of relative humidity, *J. Geophys. Res.*, *108*(D23), 8650, doi:10.1029/2003JD003405.
- Fuller, K. A. (1995), Scattering and absorption cross sections of compounded spheres. III. Spheres containing arbitrarily located spherical inhomogeneities, *J. Opt. Soc. Am.*, *12*(5), 893–904, doi:10.1364/JOSAA.12.000893.
- Heintzenberg, J. (2009), The SAMUM-1 experiment over southern Morocco: Overview and introduction, *Tellus, Ser. B*, *61*(1), 2–11, doi:10.1111/j.1600-0889.2008.00403.x.
- Howell, S. G., A. D. Clarke, Y. Shinozuka, V. Kapustin, C. S. McNaughton, B. J. Huebert, S. J. Doherty, and T. L. Anderson (2006), Influence of relative humidity upon pollution and dust during ACE-Asia: Size distributions and implications for optical properties, *J. Geophys. Res.*, *111*, D06205, doi:10.1029/2004JD005759.
- Israelevich, P. L., et al. (2003), Annual variations of physical properties of desert dust over Israel, *J. Geophys. Res.*, *108*(D13), 4381, doi:10.1029/2002JD003163.
- Kaaden, N., et al. (2009), State of mixing, shape factor, number size distribution, and hygroscopic growth of the Saharan anthropogenic and mineral dust aerosol at Tinfou, Morocco, *Tellus, Ser. B*, *61*(1), 51–63, doi:10.1111/j.1600-0889.2008.00388.x.
- Lack, D., et al. (2006), Aerosol absorption measurement using photoacoustic spectroscopy: Sensitivity, calibration, and uncertainty developments, *Aerosol Sci. Technol.*, *40*(9), 697–708, doi:10.1080/02786820600803917.
- Lack, D. A., et al. (2008), Bias in filter based aerosol light absorption measurements due to organic aerosol loading: Evidence from ambient measurements, *Aerosol Sci. Technol.*, *42*(12), 1033–1041, doi:10.1080/02786820802389277.
- Levin, Z., et al. (1996), The effects of desert particles coated with sulfate on rain formation in the eastern Mediterranean, *J. Appl. Meteorol.*, *35*(9), 1511–1523, doi:10.1175/1520-0450(1996)035<1511:TEODPC>2.0.CO;2.

- Levin, Z., A. Teller, E. Ganor, and Y. Yin (2005), On the interactions of mineral dust, sea-salt particles, and clouds: A measurement and modeling study from the Mediterranean Israeli Dust Experiment campaign, *J. Geophys. Res.*, *110*, D20202, doi:10.1029/2005JD005810.
- Lewis, K. A., et al. (2009), Reduction in biomass burning aerosol light absorption upon humidification: Roles of inorganically-induced hygroscopicity, particle collapse, and photoacoustic heat and mass transfer, *Atmos. Chem. Phys. Discuss.*, *9*(4), 15,247–15,294.
- Li-Jones, X., et al. (1998), Effect of relative humidity on light scattering by mineral dust aerosol as measured in the marine boundary layer over the tropical Atlantic Ocean, *J. Geophys. Res.*, *103*(D23), 31,113–31,121, doi:10.1029/98JD01800.
- Linke, C., et al. (2006), Optical properties and mineralogical composition of different Saharan mineral dust samples: A laboratory study, *Atmos. Chem. Phys.*, *6*(11), 3315–3323.
- McNaughton, C. S., et al. (2009), Observations of heterogeneous reactions between Asian pollution and mineral dust over the eastern North Pacific during INTEX-B, *Atmos. Chem. Phys. Discuss.*, *9*(2), 8469–8539.
- Müller, T., et al. (2009), Spectral absorption coefficients and imaginary parts of refractive indices of Saharan dust during SAMUM-1, *Tellus, Ser. B*, *61*(1), 79–95, doi:10.1111/j.1600-0889.2008.00399.x.
- Murphy, D. M. (2009), The effect of water evaporation on photoacoustic signals in transition and molecular flow, *Aerosol Sci. Technol.*, *43*(4), 356–363, doi:10.1080/02786820802657392.
- Noble, C. A., and K. A. Prather (1996), Real-time measurement of correlated size and composition profiles of individual atmospheric aerosol particles, *Environ. Sci. Technol.*, *30*(9), 2667–2680, doi:10.1021/es950669j.
- Quinn, P. K., and T. S. Bates (2005), Regional aerosol properties: Comparisons of boundary layer measurements from ACE 1, ACE 2, Aerosols99, INDOEX, ACE Asia, TARFOX, and NEAQS, *J. Geophys. Res.*, *110*, D14202, doi:10.1029/2004JD004755.
- Rosenfeld, D., and Y. Rudich (2001), Desert dust suppressing precipitation: A possible desertification feedback loop, *Proc. Natl. Acad. Sci. U. S. A.*, *98*, 5975–5980, doi:10.1073/pnas.101122798.
- Rudich, Y., O. Khersonsky, and D. Rosenfeld (2002), Treating clouds with a grain of salt, *Geophys. Res. Lett.*, *29*(22), 2060, doi:10.1029/2002GL016055.
- Schmid, O., et al. (2006), Spectral light absorption by ambient aerosols influenced by biomass burning in the Amazon Basin. I: Comparison and field calibration of absorption measurement techniques, *Atmos. Chem. Phys.*, *6*(11), 3443–3462.
- Schmidt, G. A., et al. (2006), Present-day atmospheric simulations using GISS ModelE: Comparison to in situ, satellite, and reanalysis data, *J. Clim.*, *19*(2), 153–192, doi:10.1175/JCLI3612.1.
- Shell, K. M., and R. C. J. Somerville (2007), Sensitivity of climate forcing and response to dust optical properties in an idealized model, *J. Geophys. Res.*, *112*, D03206, doi:10.1029/2006JD007198.
- Solmon, F., M. Mallet, N. Elguindi, F. Giorgi, A. Zakey, and A. Konaré (2008), Dust aerosol impact on regional precipitation over western Africa, mechanisms and sensitivity to absorption properties, *Geophys. Res. Lett.*, *35*, L24705, doi:10.1029/2008GL035900.
- Stohl, A., et al. (2005), Technical note: The Lagrangian particle dispersion model FLEXPART version 6.2, *Atmos. Chem. Phys.*, *5*(9), 2461–2474.
- Tang, I. N., and H. R. Munkjelwitz (1994), Water activities, densities, and refractive indices of aqueous sulfates and sodium nitrate droplets of atmospheric importance, *J. Geophys. Res.*, *99*(D9), 18,801–18,808, doi:10.1029/94JD01345.
- Trochkin, D., et al. (2003), Mineral aerosol particles collected in Dunuang, China, and their comparison with chemically modified particles collected over Japan, *J. Geophys. Res.*, *108*(D23), 8642, doi:10.1029/2002JD003268.
- Tucker, S. C., et al. (2009), Doppler Lidar estimation of mixing height using turbulence, shear, and aerosol profiles, *J. Atmos. Oceanic Technol.*, *26*(4), 673–688, doi:10.1175/2008JTECHA1157.1.
- Usher, C. R., et al. (2003), Reactions on mineral dust, *Chem. Rev.*, *103*(12), 4883–4939, doi:10.1021/cr020657y.
- Yang, M., et al. (2009), Attribution of aerosol light absorption to black carbon, brown carbon, and dust in China—Interpretations of atmospheric measurements during EAST-AIRE, *Atmos. Chem. Phys.*, *9*(6), 2035–2050.
- Zhang, D., Y. Iwasaka, G. Shi, J. Zang, A. Matsuki, and D. Trochkin (2003), Mixture state and size of Asian dust particles collected at southwestern Japan in spring 2000, *J. Geophys. Res.*, *108*(D24), 4760, doi:10.1029/2003JD003869.
- Zhang, R., et al. (2008), Variability in morphology, hygroscopicity, and optical properties of soot aerosols during atmospheric processing, *Proc. Natl. Acad. Sci. U. S. A.*, *105*(30), 10,291–10,296, doi:10.1073/pnas.0804860105.

T. S. Bates, D. Coffman, and P. K. Quinn, Pacific Marine Environment Laboratory, NOAA, 7600 Sand Point Way NE, Bldg. 3, Seattle, WA 98115, USA.

T. Baynard, D. A. Lack, E. Lovejoy, D. M. Murphy, A. R. Ravishankara, and S. Tucker, Earth Systems Research Laboratory, NOAA, 325 Broadway, Boulder, CO 80304, USA. (daniel.lack@noaa.gov)

D. S. Covert, Atmospheric Sciences Department, University of Washington, Box 354235, Seattle, WA 98195, USA.

P. Massoli, Aerodyne Research, Inc., 45 Manning Rd., Billerica, MA 01821, USA.

B. Sierau, Institute for Atmospheric and Climate Science, Swiss Federal Institute of Technology, Universitaetstrasse 16, CH-8092 Zurich, Switzerland.